

**APPROVED**



# Supplemental Labeling

EPA Reg. No. 241-322

## FOR USE IN PINEAPPLE FIELDS AND IN UNCULTIVATED LAND SURROUNDING PINEAPPLE FIELDS TO CONTROL PEST ANTS

SEE THE AMDRO<sup>®</sup> PRO LEAFLET LABEL FOR STATEMENTS  
OF FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

### DIRECTIONS FOR USE

This label must be in the possession of the user at the time of pesticide application.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product by any method not specified on this label.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Apply in pineapple fields and in uncultivated land surrounding pineapple fields to control pest ants.

| <b>Treatment</b>  | <b>RATE</b>                       | <b>Comments</b>  |
|---|-----------------------------------|--|
| Individual mound:<br>imported fire ants<br>native fire ants<br>bigheaded ants | 5 level tablespoons               | Distribute bait uniformly around the mound. Do not disturb the mound. Do not contaminate kitchen utensils by use or storage.   |
| Broadcast treatment:<br>imported fire ants<br>native fire ants                | 1.0 - 1.5<br>lb./acre/application | Broadcast by ground or aerial equipment (helicopter only) using disc or turbine blower type spreaders to all infested fields. Apply the first application approximately one month after planting. Apply AMDRO PRO no more than 4 applications per 12-month period with minimum intervals of 90 days. |

(see reverse side)

|  |   |  |
|--|---|--|
| Broadcast treatment:<br>bigheaded ants | 1.0 – 1.5<br>lb./acre/application<br>for moderate<br>infestations | Broadcast by ground or aerial<br>equipment (helicopter only) using disc<br>or turbine blower type spreaders to all<br>infested fields. Apply the first<br>application approximately one month<br>after planting. Apply AMDRO <sup>®</sup> PRO<br>no more than 4 applications per 12-<br>month period with minimum intervals<br>of 90 days. |
|  | 1.5 – 3.0<br>lb./acre/application<br>for heavier<br>infestations  |  |

Conduct ant surveys to monitor ant populations before and after treatments. Apply when ants are active (typically when soil temperature is greater than 60°F).

Do not exceed 6.0 lbs. of AMDRO<sup>®</sup> PRO per acre within any 12-month period.

Aerial applications are prohibited within 200 ft. of irrigation ditches, streams, ponds and public roadways, and 1000 ft. of residential areas.

Do not harvest within thirty (30) days after last application.

Do not make applications if it is likely to rain within an hour of application.

Green chop (leaves and crowns) shall not be fed to animals.

©2003 BASF Corporation  
All rights reserved

000241-00213-20031002  
NVA 2003-04-166-0213

**BASF Corporation**  
26 Davis Drive  
Research Triangle Park, NC 27709

**BASF**