DIRECTIONS FOR USE / USE LIMITATIONS: Bait MUST be placed in tamper-resistant bait stations. This bait station may be used in indoor areas accessible to children, consistent with all use restrictions and other requirements indicated on this label. DO NOT USE THIS PRODUCT OUTDOORS OR IN AREAS ACCESSIBLE TO PETS, DOMESTIC ANIMALS, OR NONTARGET WILDLIFE. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Selection of Treatment Areas: [House mice are excellent climbers and will infest all floors in a structure. Mice will only range approximately 10 m from their nesting area seeking food.] Place bait in locations mice travel and will most likely find and consume the bait. Placement along walls, in corners, by gnawed openings, in corners and concealed places, between floors or walls, or in locations where house mice or their signs have been seen. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

Remove bait station from package and place in horizontal position with side against wall, exposing entrance.

To Load:

- 1. Insert key in hole on top of mouse bait station and turn key 90°.
- 2. Lift lid and remove key.
- 3. Place 1 bait block in the square compartments.
- 4. Close lid until it snaps making certain that station is locked.

Place 2 bait stations at intervals of 2-4 m within the treatment area. If a bait placement is not disturbed after several days, move the placement approximately 1 meter. Four (4) bait stations per placement may be necessary at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

Replace bait stations if damaged or contents are consumed or contaminated. Wearing rubber gloves, collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation occurs, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations as necessary.

DOMESTIC



REFILLABLE BAIT STATION

RE-USEABLE/LOCKABLE BAIT STATION RODENTICIDE

INCLUDES LOCKING KEY

THIS BAIT STATION IS RESISTANT TO TAMPERING BY CHILDREN. USE INDOORS ONLY

GUARANTEE: Chlorophacinone 0.005%
REGISTRATION NO. 30799
PEST CONTROL PRODUCTS ACT

DO NOT ALLOW CHILDREN OR PETS TO HANDLE BAIT OR BAIT STATION READ THE LABEL BEFORE USING WARNING - CONTAINS THE ALLERGENS PEANUTS, WHEAT, SOY AND FISH



Liphatech, Inc. 3600 W. Elm Street Milwaukee, WI 53209 USA 800-351-1476

PRECAUTIONS: KEEP AWAY FROM CHILDREN, PETS AND

LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Rubber gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

FIRST AID: If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed: For all cases of human ingestion call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If pet or livestock poisoning is suspected, immediately contact a veterinarian.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Vitamin K_1 in the form of intramuscular or subcutaneous injections, or by oral ingestion is a suggested remedial treatment for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE AND DISPOSAL:

Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals. Wrap unwanted or damaged bait stations and discard in household garbage. DO NOT use damaged bait station.

RODENT DISPOSAL: Dead rodents should be securely wrapped and placed in closed containers and disposed of in the garbage. Otherwise, securely wrap carcasses and bury to a depth that will make them inaccessible to scavengers.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product. (031014)