

# SOCEPTER®

T & O | 70WDG HERBICIDE

A selective Herbicide for Use on Established Lawns of Bermudagrass, St. Augustinegrass, Centipedegrass, and Zoysiagrass.

For use on Golf Courses, Sod Farms, Commercial and Residential Turf, and for use around selected Landscape Ornamentals.

**Active Ingredient:**

Imazaquin (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-quinolinecarboxylic acid)..... 70.0%

**Inert Ingredients**..... 30.0%

**TOTAL** ..... 100.0%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION/PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**FIRST AID**

<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If In eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

**EMERGENCY INFORMATION**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.  
FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:  
For Medical Emergencies phone: .....1-888-681-4261  
For Transportation Emergencies, including spill, leak or fire, phone: CHEMTREC® .....1-800-424-9300  
For Product Use Information phone: AMVAC® .....1-888-462-6822

See inside for complete **Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

EPA Reg. No. 5481-613  
Product of U.S.A.

EPA Est. No. \_\_\_\_\_  
Net Contents: \_\_\_ pounds



AMVAC Chemical Corporation  
4100 E. Washington Blvd.  
Los Angeles, CA 90023 U.S.A.  
1-888-462-6822

---

## PRECAUTIONARY STATEMENTS

---

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Causes eye irritation. Avoid contact with skin, eyes, or clothing.  
Harmful if absorbed through skin.

#### Personal Protective Equipment (PPE):

##### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical resistant gloves.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

##### User Safety Recommendations:

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### ENVIRONMENTAL HAZARDS

**DO NOT** apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark.

**DO NOT** contaminate water by cleaning of equipment or disposal of waste. This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

---

#### DIRECTIONS FOR USE

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Observe all cautions and limitations on this label and on the labels of products used in combination with Scepter® T&O 70 WDG Herbicide. The use of Scepter T&O 70 WDG Herbicide not consistent with this label may result in plant injury. Keep containers closed to avoid spills and contamination.

**DO NOT** apply this product through any type of irrigation system.

**DO NOT** apply this product aurally.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of **12 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls.
- Chemical resistant gloves.
- Shoes plus socks.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of the product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**DO NOT** enter treated areas without protective clothing until sprays have dried.

### USE INFORMATION

Scepter® T&O 70 WDG Herbicide is a unique, versatile herbicide that will control or aid in the control of many important weeds in warm-season turfgrasses and selected ornamentals. Scepter T&O 70 WDG Herbicide may be used on established warm-season turfgrasses Bermudagrass, St. Augustinegrass, Centipedegrass and Zoysiagrass, to grounds or lawns around residential and commercial establishments, parks, golf courses, athletic fields, cemeteries, and sod farms, and to control specified weeds in ornamentals in the maintained landscape.

Scepter T&O 70 WDG Herbicide will effectively control susceptible weeds when applied during the directed application timings (pre and post).

Scepter T&O 70 WDG Herbicide-treated turfgrasses may have a compacted growth habit and formation of seedheads may be inhibited for 2-6 weeks depending on grass species and growing conditions at the time of application. Turf vigor is not impaired with this application when the rate, timing and turf species are followed according to label directions.

The activity of Scepter T&O 70 WDG Herbicide involves uptake by weed roots and/or foliage and rapid translocation to growing points. Treated plants stop growing soon after spray application. Chlorosis appears first in the newest leaves, with necrosis spreading from this point. In perennials, the herbicide is translocated into, and kills, underground storage organs. Complete death may not occur for several weeks after application in susceptible weed species. The amount of rainfall or irrigation required following application depends on existing soil moisture, density of turf and thatch or other surface vegetation, and soil type. Sufficient rainfall or irrigation of at least 0.25 to 0.5 inches is necessary for optimum herbicide activation.

### HERBICIDE COMBINATIONS

When Scepter T&O 70 WDG Herbicide is used in tank-mixture with another herbicide refer to each label for rates, methods of application, proper timing, weeds controlled, limitations and precautions. Scepter T&O 70 WDG Herbicide cannot be mixed with any product containing a label prohibition against such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### For Warm-Season Turfgrasses

Scepter T&O 70 WDG Herbicide may be tank-mixed with registered preemergence and broadleaf postemergence herbicides and with methylarsonate herbicides, such as MSMA, to control weeds not controlled by Scepter T&O 70 WDG Herbicide alone. **DO NOT** tank-mix Scepter T&O 70 WDG Herbicide with other postemergence herbicides labeled for grass weed control in turf. Scepter T&O 70 WDG Herbicide may be applied prior to or sequential to the use of all currently labeled preemergence and postemergence herbicides.

Scepter T&O 70 WDG Herbicide may be tank-mixed with urea or liquid iron for routine, turf-maintenance applications. Add Scepter T&O 70 WDG Herbicide to the spray tank first and make sure it is thoroughly mixed before adding the nutrient solutions.

### **For Landscape Ornamentals**

Scepter T&O 70 WDG Herbicide may be tank-mixed with Pendulum® 3.3 EC herbicide or Pendulum® WDG herbicide or PRE-M® 3.3 EC Turf herbicide or PRE-M 60 DG herbicide to control weeds not controlled by Scepter T&O 70 WDG Herbicide alone.

### **MIXING INSTRUCTIONS**

Fill the spray tank one-half to three-quarters full with clean water. While agitating add the required amount of Scepter T&O 70 WDG Herbicide and then fill the remainder of the tank with water. Maintain agitation while spraying to ensure a uniform spray mixture.

When tank-mixing Scepter T&O 70 WDG Herbicide with directed herbicides, add Scepter T&O 70 WDG Herbicide to the spray tank first and make sure it is thoroughly mixed before adding the tank-mix herbicide.

### **ADDITION OF SURFACTANTS**

A non-ionic surfactant at the rate of 2 pints per 100 gallons (0.25% v/v) should be added to the spray mixture. Add the non-ionic surfactant to the spray mixture after herbicide(s) have mixed. An antifoaming agent may be added to the tank, if needed.

### **SPRAYING INSTRUCTIONS**

Uniformly apply Scepter T&O 70 WDG Herbicide sprays with a properly calibrated sprayer in sufficient volume to ensure adequate coverage (20-200 gallons/acre). AVOID drift onto vegetables, flowers, ornamental shrubs and other desirable plants or injury may result. AVOID overlaps when spraying. A spray indicator dye may be added to prevent overlaps. **DO NOT** apply when wind velocity is greater than 10 mph.

### **APPLICATION RATES AND DIRECTIONS FOR USE ON WARM-SEASON TURFGRASSES**

#### **Bermudagrass, St. Augustinegrass, Centipedegrass and Zoysiagrass**

Scepter T&O 70 WDG Herbicide can be selectively used to control weeds anytime during the year **except prior to and during transition from turfgrass winter dormancy**. Application rates applied are 0.2 ounces to 0.26 ounces of Scepter T&O 70 WDG Herbicide per 1,000 sq. ft. per application, or applied at 8.6 ounces to 11.4 ounces of product per acre (or .375 lb. to 0.5 lb. ai./A) per application. Repeat applications may be necessary but do not apply a total of more than 0.75 lb. ai per acre annually. The high rate should be used where heavy weed infestations occur.

For SPOT SPRAYING with a hand-held or back-pack sprayer, mix 0.5 ounces of Scepter T&O 70 WDG Herbicide per 3 gallons of water and spray to wet. [Note: Approximately 1.5 gallons of spray solution will treat 1,000 sq. ft..

Tolerant turfgrasses metabolize Scepter 70 WDG T&O Herbicide. Best tolerance occurs under totally dormant or optimal growing conditions. When the turfgrass is growing under marginal conditions, such as low temperatures, drought stress, compaction, etc., temporary yellowing or browning can occur and recovery may take 2-4 weeks.

If product is applied where runoff is likely to occur, including embankments or steep slopes, irrigation should be applied (0.25 to 0.5 inches) to incorporate the product into the soil to prevent injury of nearby sensitive turf such as over-seeded areas, bentgrass and perennial ryegrass putting greens. When making applications adjacent to sensitive turfgrasses including bentgrass and ryegrass, care should be taken not to track the Scepter T&O 70 WDG Herbicide onto these areas.

Irrigation or rainfall of 0.25 to 0.5 inches is suggested within 1-7 days after application to wash Scepter T&O 70 WDG Herbicide into the shoot/root zone where greatest weed control activity occurs.

## CONTROL OF SUMMER WEEDS

For the postemergence control of weeds including the nutsedges<sup>1+2</sup>, dollarweed<sup>2</sup>, kyllinga<sup>1+2</sup>, field sandbur<sup>2</sup>, and ryegrass<sup>2</sup>, applications of Scepter T&O 70 WDG Herbicide should be timed following spring transition of the turf and prior to the onset of winter dormancy. This application timing would be the period of active turf and summer weed growth. **Refer to all turfgrass precautions prior to use.**

## CONTROL OF WINTER WEEDS

For the control of weeds including wild garlic<sup>3</sup>, wild onion<sup>3</sup>, henbit, chickweed, lawn burweed, etc., applications should be timed to follow the first frost. Scepter T&O 70 WDG Herbicide may be applied prior to or soon after emergence of the weeds listed with the exception of wild onion and wild garlic which must be emerged at application time for control. Control of emerged weeds may take several weeks. Applications made prior to the first frost or later during the dormant period (inactive weed growth) may result in reduced weed control. Applications must not be made less than six weeks prior to the initiation of spring transition. **Refer to all turfgrass precautions prior to use.**

## WEED SPECIES CONTROLLED

Common Name	Genus species	Annual/Biennial/ Perennial
<b>GRASSES CONTROLLED</b>		
Field Sandbur	<i>Cenchrus incertus</i>	Annual/short-lived perennial
Perennial Ryegrass <sup>2</sup> (Over-seeded)	<i>Lolium perenne</i>	Perennial/winter annual
<b>SEDGES CONTROLLED</b>		
Globe Sedge <sup>1+2</sup>	<i>Cyperus globosus</i>	Perennial
Green Kyllinga <sup>1+2</sup>	<i>Kyllinga brevifolia</i>	Perennial
Rice Flatsedge <sup>1+2</sup>	<i>Cyperus iria</i>	Annual
Purple Nutsedge <sup>1+2</sup>	<i>Cyperus rotundus</i>	Perennial
Yellow Nutsedge <sup>1+2</sup>	<i>Cyperus esculentus</i>	Perennial
Wild Garlic <sup>3</sup>	<i>Allium vineale</i>	Winter perennial
Wild Onion <sup>3</sup>	<i>Allium canadense</i>	Winter perennial
<b>BROADLEAVES CONTROLLED</b>		
Black Medic	<i>Medicago lupulina</i>	Annual
Buttercup	<i>Ranunculus Parviflorus</i>	Winter annual
Carpetweed	<i>Mollugo verticillata</i>	Annual
Chickweed Common	<i>Stellaria media</i>	Winter annual
Cutleaf Evening Primrose	<i>Oenothera lanceolate</i>	Winter annual
Dollarweed <sup>2</sup>	<i>Hydrocotyle umbellate</i>	Perennial
Dovestool	<i>Geranium molle</i>	Winter annual
Eclipta	<i>Eclipta prostrata</i>	Annual
Geranium Carolina	<i>Geranium carolinianum</i>	Winter annual
Hairy Bittercress	<i>Cardamine hirsuta</i>	Winter annual
Henbit	<i>Lamium amplexicaule</i>	Winter annual
Knawel	<i>Scleranthus annuus</i>	Winter annual
Lawn Burweed	<i>Soliva pterosperma</i>	Winter annual
Mouseear Chickweed	<i>Cerastium vulgatum</i>	Winter annual
Parsley-piert	<i>Alchemilla arvensis</i>	Winter annual
Pigweed species	<i>Amaranthus sp.</i>	Annual
Purple Deadnettle	<i>Lamium purpureum</i>	Winter annual
Red Sorrel	<i>Rumex acetosella</i>	Perennial
White Clover	<i>Trifolium repens</i>	Annual
<b>Scepter T&amp;O 70 WDG Herbicide will aid in the control of and reduce competition from:</b>		
Annual Bluegrass	<i>Poa annua</i>	Winter annual

Bahiagrass	<i>Paspalum notatum</i>	Perennial
Crabgrass species	<i>Digitaria sp.</i>	Annual
Tall Fescue	<i>Festuca arundinacea</i>	Perennial
Annual Sedge <sup>1,2</sup>	<i>Cyperus compressus</i>	Annual
Cudweed, Purple	<i>Gnaphalium purpureum</i>	Annual/biennial
Dandelion	<i>Taraxacum officinale</i>	Winter annual
Violets	<i>Viola sp.</i>	Winter annual/perennial
Virginia Buttonweed	<i>Diodia virginiana</i>	Perennial

<sup>1</sup> For control of kyllinga and more rapid and improved control of the sedges the addition of MSMA (2 lb. ai/A) is suggested in methylarsonate tolerant turfgrasses.

<sup>2</sup> A repeat application may be required for extended postemergence control of weeds including dollarweed, Kyllinga, the nutsedges, and perennial ryegrass.

<sup>3</sup> Mowing of wild onion/garlic 1-2 weeks after treatment is recommended for improved appearance and initial control.

### SPECIAL RESTRICTIONS ON WARM SEASON TURFGRASSES

- USE ONLY on well-established non-stressed turfgrass with a dense and uniform stand. TEMPORARY YELLOWING may occur.
- **DO NOT** apply just prior to or during spring transition or during periods of very slow turf growth as severe discoloration could occur.
- **DO NOT** use on golf or bowling greens.
- **DO NOT** use on unlabeled turfgrass species, including dichondra.
- **DO NOT** APPLY Scepter T&O 70 WDG Herbicide to St. Augustinegrass for winter weed control.
- **DO NOT** mow St. Augustinegrass until 48 hours after application.
- **DO NOT** use on tall fescue or mixed stands of tall fescue and Bermudagrass when tall fescue is a desirable turfgrass. Scepter T&O 70 WDG Herbicide may injure tall fescue.
- **DO NOT** use on winter over-seeded turfgrasses such as ryegrass, bentgrass or *Poa trivialis* when those species are desirable.
- **DO NOT** reseed or winter overseed or sprig turfgrass for 1.5 months (six weeks) after treatment.
- **DO NOT** apply to newly planted, sprigged or sodded turfgrass.
- **DO NOT** apply within 30 days of harvesting sod.
- **DO NOT** graze or feed clippings of treated turfgrasses to animals.
- **DO NOT** use treated grass clippings as mulch in landscape beds or around vegetables in the garden or fruit trees, vines or berries in the landscape areas.

### DIRECTIONS FOR USE FOR LANDSCAPE ORNAMENTALS

Scepter T&O 70 WDG Herbicide can be safely applied at the labeled use rates around and over the top of those plants listed below for the control of weeds listed on this label. Applications can be made safely around the drip zone of deciduous and conifer trees in the landscape where established stands of turfgrass are present. To make applications to ornamentals or trees in mulched landscape beds consult the list of tolerant species. Temporary growth suppression may be observed on some treated species.

Application rates applied are 0.2 ounces to 0.26 ounces of Scepter T&O 70 WDG Herbicide per 1,000 sq. ft. per application. For SPOT SPRAYING with a hand-held or back-pack sprayer, mix 0.5 oz. of Scepter T&O 70 WDG Herbicide per 3 gallons of water and spray to wet. [Note: Approximately 1.5 gallons of spray solution will treat 1,000 sq. ft.]

Scepter T&O 70 WDG Herbicide is for landscape ornamental and turfgrass use only.

**DO NOT** APPLY this product around food producing trees or vines, berry bushes or in vegetable gardens.

**DO NOT** APPLY this product in field- or container-grown ornamental nurseries, in nursery liner, seedling or tree nurseries.

## TOLERANT LANDSCAPE ORNAMENTAL SPECIES

Common Name	Genus species 'Cultivar or Variety'
<b>SHRUBS</b>	
Indian Hawthorne	<i>Rhaphiolepis indica</i>
Dwarf Yaupon Holly	<i>Ilex vomitoria</i> 'Schilling's Dwarf'
Blue Pfitzer Juniper	<i>Juniperus chinensis</i> 'Pfitzeriana Glauca'
Helleri Holly	<i>Ilex crenata</i> 'Helleri'
Red Tip Photinia	<i>Photinia X fraseri</i>
Yucca	<i>Yucca pendula</i>
Wax Myrtle	<i>Myrica cerifera</i>
Burford Holly	<i>Ilex cornuta</i> 'Burfordii'
Gardenia	<i>Gardenia jasminoides</i>
'Miami Supreme' Flame of the Woods*	<i>Ixora coccinea</i> *
Confederate Jasmine	<i>Trachelospermum jasminoides</i>
Parsons Juniper	<i>Juniperus squamata expansa</i> 'Parsonii'
Blue Pacific Juniper	<i>Juniperus conferta</i>
Blue Rug Juniper	<i>Juniperus horizontalis</i> 'Wiltonii'
<b>TREES</b>	
Red Crape Myrtle	<i>Lagerstroemia indica</i>
<b>GROUNDCOVERS</b>	
Asiatic Jasmine	<i>Trachelospermum asiaticum</i> 'Minima'
Giant Liriope	<i>Liriope muscari</i> 'Evergreen Giant'
Variiegated Liriope	<i>Liriope muscari</i> 'Silvery Sunproof'
Mondo Grass	<i>Ophiopogon japonicus</i>
Pachysandra	<i>Pachysandra terminalis</i>
<b>PERENNIALS</b>	
Hosta 'Lancifolia'	<i>Hosta</i> sp.
Society Garlic*	<i>Tulbaghia violacea</i> *

\* This plant has not been tested for tolerance to Pendulum® or PRE-M® herbicides. **DO NOT** tank mix Scepter® T&O 70 WDG Herbicide with pendimethalin herbicides and apply to this plant species.

### SPECIAL PRECAUTIONS IN LANDSCAPE ORNAMENTALS

- USE ONLY on well-established, non-stressed ornamentals **listed on this label** or injury may result. **DO NOT** apply around annual bedding plants.
- **APPLY** Scepter T&O 70 WDG Herbicide only to the listed plants or severe injury may result. The following ornamentals are known to be severely injured by Scepter T&O 70 WDG Herbicide: Azalea, Viburnum, Pieris, Abelia, and *Ligustrum*.
- **DO NOT APPLY** this product to bareground areas of the landscape where new ornamental plantings (including annual bedding plants) will be installed within the year.

### USES WITH OTHER PRODUCTS (TANK-MIXES)

If to the extent consistent with applicable law this product is used in combination with any other product except as specifically directed in writing by AMVAC then AMVAC shall have no liability for any loss, damage, or injury arising out of its use in any such combination not so specifically directed. If used in a combination directed by AMVAC the liability of AMVAC shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the AMVAC product in such combination use, and in any event shall be limited to return of the amount of the purchase price of the product.

### STORAGE AND DISPOSAL

**Prohibitions:** DO NOT contaminate water, food, or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### CONTAINER DISPOSAL

**Nonrefillable Container. DO NOT reuse or refill this container.** Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**Triple rinse containers small enough to shake (capacity ≤ 50 pounds) as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Triple rinse containers too large to shake (capacity >50 pounds) as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse as follows:** Empty the remaining contents into application equipment or mix tank. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

---

### LIMITED WARRANTY AND DISCLAIMER

---

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use, subject to the inherent risks referred to herein, when it is used in accordance with such directions; and (c) that the directions, warnings, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants and residues on food crops, and upon reports of field experience. Tests have not been made on all varieties of food crops and plants, or in all states or under all conditions.

**THERE ARE NO EXPRESS WARRANTIES OTHER THAN THOSE SET FORTH HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE, TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OF FITNESS FOR A PARTICULAR PURPOSE, OR ANY WARRANTY OF QUALITY OR PERFORMANCE. THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.**



**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

AMVAC offers this product, and Buyer accepts it, subject to the foregoing Limited Warranty which may be varied only by agreement in writing signed by an authorized representative of AMVAC.

© 2017 AMVAC Chemical Corporation. A wholly owned subsidiary of American Vanguard Corporation. All Rights Reserved. American Vanguard, AMVAC, and the AMVAC logo are trademarks owned by AMVAC Chemical Corporation.

CHEMTREC is a service mark owned by the American Chemistry Council, Inc.

Scepter is a trademark owned by BASF and used under license.

Pendulum is a trademark owned by BASF.

*PRE-M* is a trademark owned by Lesco, Inc.

AMVAC Chemical Corporation  
4100 E. Washington Blvd.  
Los Angeles, CA 90023  
1-888-462-6822