### TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian at

### NOTE TO VETERINARIAN

Anticoagulant Bromadiolone: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K1. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

#### SEE LABEL ON OUTER PACKAGING FOR ADDITIONAL PRECAUTIONARY INFORMATION

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. See outer package label for complete DIRECTIONS FOR USE including USE RESTRICTIONS, and APPLICATION DIRECTIONS. It is illegal to sell this product unless it is accompanied by a complete set of its accepted labeling.

INDIVIDUAL SALE IS PROHIBITED BY LAW

>2008-9

# CONTRAC® RODENTIGIDE READY-TO-USE PLACE PACS

### KILLS RATS, MICE & MEADOW VOLES\*

ACTIVE INGREDIENT:

## KEEP OUT OF REACH OF CHILDREN CAUTION

See back panels for First Aid and additional precautionary statements.

NET WT: **1.5 OZ** (43 g)

EPA REG. NO. 12455-76
EPA EST. NOs. 12455-WI-1<sup>k</sup>, 12455-WI-2<sup>p</sup>, 12455-WI-3<sup>w</sup>
Superscript is first letter of the lot number.

\*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, Whitethroated woodrat, Southern plains woodrat, and Mexican woodrat

Manufactured by:



Madison, WI 53704 U.S.A. www.belllabs.com

FIRST AID
HAVE LABEL WITH
YOU WHEN OBTAINING
TREATMENT ADVICE

### If Swallowed

- Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.
- Have a person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.

#### If in Eyes

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice

### NOTE TO PHYSICIAN

If swallowed or absorbed through the skin, this material may reduce the dotting ability of the blood and cause bleeding. If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

102618/09-15