



ENVIRO PROTECTION INDUSTRIES COMPANY, INC.


GRANULAR REPELLENT
PROFESSIONAL



Available at


UNIVAR[®]

Seasonal Maps by Animal



DEER ACTIVITY

Deer are found throughout the US except in the driest deserts. The two major species are White-tailed Deer, with 14 subspecies and Mule Deer with one major subspecies - the Black-tailed Deer found in the Pacific Northwest and coastal areas of Northern California, Oregon and Washington. Deer species overlap in range in the western US.



FOCUS ON PEAK APPLICATION TIMES

Increased activity ranges are ideal for repeat customer sales. Specific winter applications are available for heavy snowfall regions.



RABBIT ACTIVITY

There are 12 species of rabbits in the United States, with Cottontails being the most widespread. Due to their great adaptability rabbits thrive in many different habitats, ranging from rural fields to suburban and urban backyards. Tell-tale signs of rabbit activity in a garden or yard are: distinctive tracks, pea-sized round droppings, gnaw marks 1 to 2 feet above ground on trees and shrubs, and plant stems that are cleanly cut through. Rabbits will eat a wide variety of green vegetation, including many of the flowering plants, ornamental trees, fruits, and vegetables that gardeners prize most.



FOCUS ON PEAK APPLICATION TIMES

Rabbits are active all year long. Protect spring green growth with early application and reapply during summer. Winter damage to shrubs can be prevented by applying on top of snow & hanging small sacks on target vegetation.