

SPECIALLY FORMULATED FOR MICE







PRODUCT FEATURES:

- Specially formulated to appeal to the feeding preferences of mice
 - Superior palatability and effectiveness for difficult mouse infestations
- ▶ Lower cost per placement
 - 15gm blox provide maximum baiting flexibility and the smallest legal block placement in the industry
- Lower toxicity, lower chemical footprint
 - Our 0.0025% formula is the lowest toxicity Brodifacoum bait available
 - Reduces possibility of secondary poisoning
- Mold and moisture resistant
 - Incorporates an advanced preservative system
- ▶ Full rodent label for application flexibility
 - While optimized for mice, Formus is also a very capable rodenticide for rats. Thus, it is registered and labeled for both mice and rats





Bell More Than Meets The Eye

Madison, Wisconsin 53704 USA | Ph: (608) 241-0202 | Fax: (608) 241-9631

www.belllabs.com

Formus All-Weather Blox

DIRECTIONS FOR USE

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only It is a violation of Federal law to use this product in a manner inconsistent with its labeling. for the sites, pests, and application methods described on this label. IMPORTANT: Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent

- Store unused product out of reach of children and pets.
- Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism
- Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

Note: Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, non-target mammals, or birds may access the bait.

in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined vehicles (ships, trains, aircraft), docks and port of terminal buildings and related structures around and USE RESTRICTIONS: This product can only be used to control Norway rats, roof rats and house mice commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport above, is prohibited. This product must not be applied directly to food or feed crops.

Burrow baiting with Formus All-Weather Blox is prohibited.

contact with food. When used in USDA inspected facilities, this product must be applied in Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct tamper-resistant bait stations. Do not broadcast bait

Do not sell this product in individual containers holding less than 16 pounds of bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and concealed places, between floors and walls, or in locations where rats or mice or their signs have been consume the bait. Generally, these areas are along walls, by gnawed openings, in corners and seen. Protect bait from rain and snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

RATS: Place 7 to 30 blocks at intervals of approximately 15 to 30 feet in appropriate locations in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until there no longer are signs of rats feeding on bait. For use in sewers, thread wire through blocks and securely attach to a stationary structure such as the bottom step of a manhole ladder or a sewer grate. Allow just enough wire for the blocks to rest on manhole benching. If benching is not present, suspend blocks a

Moisture Resistant • Mold Resistant

Kills Warfarin Resistant Norway Rats And House Mice

Norway rats and house mice may consume a lethal dose in one night's feeding with first dead rodents appearing four or five days after feeding begins.

ACTIVE INGREDIENT: Brodifacoum (CAS #56073-10-0): 0.0025%
OTHER INGREDIENTS: 99.9975%
**Charleine Denastration Renzorate TOTAL 100.00000% *Contains Denatonium Benzoate

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panels for additional precautionary statements.

HAVE	HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE
If Swallowed	If Swallowed • Call a poison control center, doctor, or 1-877-854-2494 immediately for
	treatment advice.
	 Have a person sip a glass of water if able to swallow.
	 Do not induce vomiting unless told to do so by the poison control center
	or doctor.
If in Eyes	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
•	 Remove contact lenses, if present, after the first 5 minutes, then continue
	rinsing eye.
	 Call a poison control center, doctor, or 1-877-854-2494 immediately for
	treatment advice.

NOTE TO PHYSICIAN

If swallowed, this material may reduce the dotting ability of the blood and cause bleeding. If ingested, administer Vitamin K₁ intramuscularly or orally. Vitamin K₁ is antidotal at doses of 10-20 mg (**not** mg/kg). Repeat as necessary based on monitoring of prothrombin times.

If animal eats bait, call veterinarian or 1-877-854-2494 at once. TREATMENT FOR PET POISONING NOTE TO VETERINARIAN

Anticoagulant Brodifacoum: For animals ingesting balt and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K₁ at doses of 2-5 mg/kg. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

NET WEIGHT: 18 lbs (8.2 kg)

EPA REG. NO. 12455-108

Manufactured by:

3699 Kirsman Blvd.
Addison, WI 53704 U.S.A. www.bellabs.com

EPA EST NO 12455-WI-1

APPLICATION DIRECTIONS (Continued from other panel) **DIRECTIONS FOR USE** (Continued from other panel)

few inches above the high water mark or secure blocks on a board supported by opposing steps of the ladder. Securing blocks in this manner may inhibit removal by rats or water. Use at least 20 blocks per manhole. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease. MICE: Place 1 block per placement. Space placements at 8 to 12 foot intervals in infested areas. Two blocks may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a FOLLOW-UP: Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation, limit sources of rodent food, continuous source of infestation is present, establish permanent bait stations and replenish as needed.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Keep away from children, domestic animals and pets. Do not get in eyes.

All handlers (including applicators) must wear shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

User Safety Requirements

As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. gum, using tobacco or using the toilet and change into clean clothing.

This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment

STORAGE AND DISPOSAL

Pesticide Storage: Store only in original container in a cool, dry place inaccessible to children and Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be placed in trash or delivered pets. Keep containers closed and away from other chemicals.

Container Handling: Nonrefillable container. Do not reuse or refill this container. to an approved waste disposal facility.

[Plastic:] Offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by ncineration. Do not burn, unless allowed by state and local ordinances.

implied, concerning the use or this material when such use and/or handling is contrary to label instructions.

Use and/or handling of this material when such use and/or handling is contrary to label instructions. WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of