PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Hazards to Humans and Domestic Animas
Keep away from humans, domestic animals and pets, Harmful if swallowed, absorbed through the skin, inhabd, or in eyes, Do not breathe dust, Avoid contamination of feed or foodstuffs, Avoid contact with acids. Wash all utensits, spoons, or measuring devices thoroughly after using.

Personal Protective Equipment (PPE)

All handlers, including anyone who applies bait and retrieves carcasses and unused bait, must wear waterproo

groves.
Follow manufacturers instructions for cleaning/maintaining PPE, such as gloves. If no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry, Wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS This product is toxic to wildlife and fish. Birds and other mildfle feeding on treated bair may be killed, Do not apply where runoff is likely to occur. Do not apply where runoff is likely to occur. Do not apply where product directly to water, or to areas where surface water is present or to intertibal areas below the mean injoin water mark.

ENDANGERED SPECIES CONSIDERATIONS

Whooping Crane (Grus americana). Do not use this product during the period 30 days prior to arrival and 30 days after the whooping crane leaves known crane habitat.

Pyomy rabbit (Brachylagus idahoensis). Do not use this product within one mile of alfatta fields

The property of the property o Gunnison prairie dog town less than 200 acres in size; c.) has been appropriately surveyed, using methods acceptable to the U.S. Fish and Wildfie Service, and found by the FIVS not to be a suitable site for ferret reintroductions; or, d.) has been exempt by the FIVS from the survey requirement.

UTAH: Utah Prairie Dog (Cynomys parvidens). Do not use this product in the range of the Utah Prairie

CALECRNIA: Salt march harvest mouse (Reithmolontomic reviventric). Do not use this product within Octarionals: Sat intal naves intous present personal content of the content of th the Sacramento River and the San Joaquin River.

Morro Bay Kangaroo Rats (Dipodomys heermann). Do not use this product within 2 1/2 miles of Bay Wood Park which is located on Morro Bay.

Preson Cangaroo Rat (Dipodrnys nitratoides exilis). Follow the Interim Measures for Protecting Endangered Species in the California Endangered Species Bulletin for the following California counties: Fresno, Kings, Madera and Merced,

Stephan's Kangaron Bat (Diporforms stephensi). Follow the Interim Measures for Protecting Endangerer Species in the California Endangered Species Bulletin for the following California counties: Sa Jernamino, San Diego and Riverside

en lations, Sail pregular intersione. Leutian Canada Goose (Branta canadensis leucopareia). Do not use this product from October to March the Federally closed goose hunting areas in the State of California. See current Duck and Goose

Giant Kangaroo Rat (Dipodomys ingens). Follow the Interim Measures for Protecting Endangered Species in the California Endangered Species Butetin for the following California counties: Fresno, Kerr Kings, Merced, Monterey, San Benito, San Luis Obispo, Santa Barbara and Tulare.

Tipton Kangaroo Rat (Dipodomys, nitratioides nitratioides). Follow the Interim Measures for Protecting Endangered Species in the California Endangered Species Bulletin for the following California counties

San Bernardino Kangaroo Rat (Dipodo*mys merriami parvus*). Follow the Interim Measures for Protecting Endangered Species in the California Endangered Species Bulletin in the following California countie

San Bernardino and Riverside sair berination airu rivevair. PEXAS: Atheiste's greater praire chicken (Tymparuchus cupido attwateri). Do not use this product in the present occupied range of the Athwater's praire chicken in the following Texas counties: Aransas. Austin, Barazaria, Colonado, DeWitt, Fort Bend, Galveston, Codal, Fariris, Relligio, Victori, Walter and Whatron, PUERTO RICO: Yellow-shouldered Black Bird (Agelaius xamthomus) and Puerto Rican Plain Pigeon (Columba inornata wetmorel). This product must not be used in the following areas except in tamper probait boxes: within 10 kilometers of Central Aquirre, Lago Cidra, Celba, San German: 15 kilometers of L Esperanza, south of Highway 2 from city of Mayaguez to the city of Ponce and all Mona Island.

Experience, South of Highway 2 from city of Magazar to the oily of Fonce and all Mochal Bands. MOTICE: It is a foother lost own per period from a morter but read in the death of a member of an endranger species. The use of this product may pose a hazard to Friedrang Medical and endough and a member of an endangered species. Supposed endourses by product endough expecies species Modified species and product may not all EMS Endangered Species Modified. Protein Medical species from could be specied from the country and the man of the Modified Species Modified species from the Modified Species Modified from the Modified Species Modified Species Modified from the Modified Species M

DIRECTIONS FOR USE

If is a violation of Federal law to use this product in a manner inconsistent with its labeling,

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Do not apply to any site, against any pest, or by any method of application not stated on this label.

Use Restrictions (for all use patterns): Do not apply this product in a way that will contact workers or other

persons, either directly or through drift. Only protected handlers may be in the area during application. Keep all other persons out of the treated area during this application. LIMITATIONS The following Limitations apply to all agricultural areas and use sites listed on this label: This product must not be applied on roads, near residential areas, or over water. Do not broadcast over growing crops other

than alfalfa, barley, beans (dry), potatoes, sugar beets, sugarcane and wheat. Apply product on warm clear days, Do not graze animals in treated areas. HISE RESTRICTIONS: For control of California Ground Sources (Spermontilus hearbest) on noncron

rights of way adjacent to canal and ditch banks and other noncrop borders during late spring and early summer This product must not be applied on roads, near residential areas, over water or where plants are grown for foc

PREBATING: Prebailing with 6 lbs. of untreated steamed crimped oats per acre one or two days prior to using ZP® RODENT OAT BATT AG™ is likely to enhance bait acceptance by California ground squirrels, BAITING: Treat only once during treatment period. Broadcast bait in a 10-15 foot swath along right-of-way, using

hand or ground-driven dispensing devices not to exceed 6 lbs, per acre. Bait may be applied only to the canal right-of-way between the base of the levee and adjacent properties. The canal and levee may not be treated. Discose of excess bait from application equipment by buriel.

RANGELAND AND ADJACENT NONCROP AREAS

USE RESTRICTIONS: For control of black-tailed prairie dogs (Cynomys (udovicianus), white-tailed prairie dogs (C. leucurus) and Gunnison (Zun) prairie dogs (C. gunnison) on rangetand in the western United States (North Dakota, South Dekota, Nebraska, Kansas, Oldahoma, Texas, New Mexico, Arizona, Colorado, Montana, Idaho

PREBAITING: Prebait with 4 grams (one teaspoon) of untreated steamed crimped oats per mound, one or two days prior to balting, to increase acceptance of treated balts by prairie dogs,

BATTING: Apply ball only after all or most of prebail is eaten and only to areas where untreated balt was consumed. Establish observation period during prebailing. Apply ball by hand as a six-inch ball spot on edge of 621.3.

each mound or in adjacent feeding areas. Application rate should not exceed 4 grams (one teaspoon) per bait

KANGAROO RATS

HISE RESTRICTIONS: For control of the Ord's kangaron rate (Dinodomus ordi), hanner tailed kangaron rate (D USE RESTRICTIONS: For control or the vitro singstron rats (uppromise oran), cohiner-series unarganitor lats (up, specifically), and merism's leangarcor arts (0, merism) in nagleand vegetation and noncrop areas. Do not apply this product within the occupied habitats of threatened or endangered species or subspecies of kangarcor arts. HAND BATTING: Place one tablespoon (12 grams) of bail at 2 locations in feeding runs near the entrances to

MACADAMIA NUT ORCHARDS AND ADJACENT NONCROP AREAS HSF RESTRICTIONS: For control of roof rats (Rattus rattus) in maca

teaspoons (4-6 grams) of bait in bait stations at the base of each tree and in each station in trees. Uneaten bait must be removed from trees prior to shaker harvest or from ground prior to mechanized harvest. Orchards may e treated up to 30 days prior to a harvesting round.

BURROW TREATMENT: Place 1.2 teaspoons (4-8 grams) of hait in each burrow. No preharvest interval is

BUHNOW INEALMENT HEAD 12 WAS SPORTS (HE grams) or ball in each output, we premierate in requiring, provided nats are not retrieved from burnows, BROADCAST BATTING: Broadcast at the rate of five (5) pounds of ball per are in orchard using a ground device or by hand. Averait may be used to here adjacent concerp areas if recommended and supervised by the Hamail Dept. of Health. The number of applications shall not exceed four (4) per year. The total amount of bait applied shall not exceed 20 pounds per acre per year.

ORCHARDS AND GROVES

VOLES AND WHITE-FOOTED MICE USE RESTRICTIONS: For control of meadow voles, prairie voles, pine voles, California voles, long-tailed voles, Oregon voles, mountain votes, Tomsend's votes (Microtus spp.) and white-looted mice in orchards and groves.

Organ Vest, mountain vides, limited vides, incurrent vides, limited vides from the mountain vides, limited vides from the control specific vides of the control vides of the v

TRAILBUILDER: Set equipment to drop one teaspoon quantity (4 grams) of bait at 4-5 foot intervals in the rificial trail, made by the machine, just inside the drop line on both sides of the trees. Apply bait at the rate of

gloves during hand-broadcast operations. Concentrate bailing in the areas with the heaviest vegetative cover. Do not broadcast on bare ground or growing crops. Apply bail at rates from 6-10 lbs. per treated acre. AERIAL APPLICATION: Broadcast balt at a rate of 6-10 lbs, per acre. Do not apply on bare ground or growing

USE RESTRICTIONS: For control of California Ground Sourcels (Spermonhius beaches) and Richardson's Ground Squirrels (Spermophilus richardi) in orchards and groves. Apply bait after harvest while the orchard and

govers in a nunceasing insiste.

PREBATING: Prebating with 6 lbs, of untreated steamed crimped oats per acre one or two days prior to using ZP® RODENT OAT BAIT AG** is likely to enhance bait acceptance by California and Richardson's ground squirrels. HANDBAITING: Apply one teaspoon (4 grams) of bait on the ground near active burrows.

BAITING: Treat only once during treatment period. Broadcast bait using hand or ground-driven dis devices not to exceed 6 lbs, per acre, Wear rubber gloves during hand-broadcast operations. Dis excess bait from application equipment by buriel.

POCKET GOPHERS

USE RESTRICTIONS: For control of plains pocket gophers (Geomys bursarius), southeastern pocket gopher Use the INC-IUMS: Inc commot or pairs possed operate (sectings oursalized, southeastern possed operate (c) printed), selectioned possed applies (Flappageagemes calanage) and posted oppines (Thomamum spp) in orchards and growes. Bait must be applied directly into posted popher's underground burrow systems or into underground burnels created by burrow-builder machines, Do not apply bait above ground, HAND BATIMIC Follow the directions indicated below to beate and treat posted gothers' burrow systems.

Deciding Underground Figurings: Use a cooking size of the problem of the problem

Applying Bait to Runways: Using a long-handled spoon or mechanical probe, drop one teaspool each baited point, Cover the opening with sod, a rock, or soil to exclude light. Take care not to cover bait with soil. Depending upon mound density, make two to five bait placements per burrow system, Recover and bury all solled bait. Applied in this manner, one pound of bait will treat one to eight acres, depending upon the number of packet appher burrow systems present.

BURROW BUILDER: Follow manufacturer's instructions for the type of equipment used. Calibrate equi-drop a teaspoon quantity (4 grams) of bait at 4-5 foot intervals in the artificial burrow made by the machin at a rate of 2-3 bs, of bait per acre, Prok up and bury at spilled bair.

SUGARCANE

SUGARCANE sugarcane netss.

PREBATTING: Prebaiting with three pounds of untreated steamed crimped cats per acre one or two weeks prior

BAITING: Broadcast bait by aircraft, ground-driven devices, or by hand at the rate of 5 lbs, of bait per acre. Wear

rubber gloves during hand-broadcast operations. The number of applications shall not exceed 4 per crop. The total application shall not exceed 20 lbs, per acre. Do not apply within 30 days of harvest

NURSERY STOCK CONFERNMENT AS TREE POR ARCOTTONWOOD

USE RESTRICTIONS: For control of meadow votes, prairie votes, pine votes, California votes, Oregon votes, intain voles, and Townsend's voles (Microtus soc.) in nursery stock, and coniferiChristmas treood, ornamental, and non-bearing fruit tree plantations. Do not apply by air.

popularizonmologi ornamenta, ai no non-teaming ruin tire plantations. Lo no apply of six in BROADCAST BATING: Under infection tuneery stock, conferior/strimss trees, popularizottomicod trees ornamiental trees, or non-bearing ruint trees, broadcast bati everify by cyction seeder or by hand. Wear ruibbe glowere during hand-broadcast operations. Concentrate in areas with the heariest vegetative cover. Do no broadcast on bare ground, Apply at a rate of 6-10 bs, per acre.

HAND BAITING: Near the base of infested nursery stock, conifer/Christmas trees, poptar/cottonwood name benines: need line case or illnessed inasery stock, contendentialistics lines, populationalismos, trees, cramental fases, or nor-bearing from trees, place tession for quantifies (4 grams) of ball at 24-1 locations, either on surface trails or at the mouth of holes leading to underground burrow systems. Do not allow ball to be exposed on bare ground, Do not disturb the runnay system, Salt at a rate of 2-3 lbs, per acre of infested stock, GRAPE VINEYARDS

USE RESTRICTIONS: For control of deer mice (Peromyscus maniculatus), white-footed mice (P. leucopus), oldfield mice (P polionotus, except for threatened or endangered subspecies), meadow voles, prairie voles and pine voles (Microtus spp.), and meadow jumping mice (Zapus hudsonius) in grape vineyards. Do not use this product in areas of Alabama and Florida that are occurried by threatened or endangered subspecies of the

BROADCAST BAITING: Broadcast bait evenly on the ground between the rows by cyclone seeder or by hand. Wear rubber gloves during hand-broadcast operations. Apply bait at a rate of 6-10 bs, per acre. Do not apply

CROPLAND, RANGELAND, AND NONCROP AREAS

RESTRICTED USE PESTICIDE

DUE TO HAZARDS TO NONTARGET SPECIES

retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification

ZP RODENT AG

OAT GRAIN BAIT FORMULATION

FOR CONTROL OF GROUND SQUIRRELS PRAIRIE DOGS RATS VOLES MICE.

ACTIVE INGREDIENT:		
Zinc Phosphide: (CAS#	1314-84-7):	2%
INERT INGREDIENTS: .		98%
	TOTAL	100%

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

Have label with you when seeking treatment advice, including 1-877-854-2494

Any persons applying zinc phosphide products and experiencing signs and symptoms such as nausea, abdominal pain, tightness in the chest, or weakness, should be seen by a physician immediately.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice or transport patient to the nearest hospital.

Do not drink water Do not administer anything by mouth or induce vomiting unless told

to do so by the poison control center or doctor. IF ON SKIN OR CLOTHING:

Take off contaminated clothing.

Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor immediately for treatment advice.

IF INHALED: Move person to fresh air

If person is not breathing, call an ambulance, then give artificial respiration, preferably mouth to mouth if possible

Call a poison control center or doctor immediately for treatment advice IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15-20 minutes Remove contact lenses, if present, after the first 5 minutes, then

continue rinsing eve. Call a poison control center or doctor immediately for treatment advice.

See side panels for additional precautionary statements.

NET WEIGHT: 45 lbs (20.4 kg)



SQUIRREL OATS

USE RESTRICTIONS: For control of plains pocket pophers (Georges bursarius), southeastern pocket gophers of 6 lbs. (0.12 lb, active) per acre. Wear rubber gloves during hand-broadcast operations. No more than one use mean-minutes run control of paries poose gapteres (seemine bussianus, southeastern pooset gapteres (or, predits), puller bussianus, pooset gapteres (seemine bussianus, southeastern pooset gapteres cropland (cifuding) attitute, pooset gapteres, referentation areas. Orinitainas trees, conferiopolaricotimosed particulares and conferentation pooset gapteres and pooset pool pooset gapteres and pooset gapteres and poolet gapteres and poolet

HAND BAITING: Follow the directions indicated below to locate and treat pocket gophers' burrow systems. <u>Locating Underground Runways:</u> Use a specialty designed popher probe, a metal rod, a strong smooth stick, or other suitable implement to probe the ground 10 to 15 inches away from fan-shaped gopher mounds. Begin probing on the fat side of the fan. When the main runway has been entered, the probe will drop about 2 inches prouning on the fact side of the fact. When the main formary has been entered, the probe will drop about 2 mones due to decreased resistance from soil. <u>Applying Bail to Riumags</u>: Using a long-handled spoon or mechanical probe, drop one teaspoon (4 grams) of bail

into the main runway at each baited point. Cover the opening with sod, a rock, or soil to exclude light. Take care not to over bait with sol. Depending upon mound density, make two to five bait placements per burrow system. Recover and burval soil debait, Accided in this manner, one pound botal will treat one to eight acres, depending non the number of packet ganher burrow systems present.

upon me numeer or poows gopiner burrow systems present.

BURROW BUILDER: Follow manufacturer's instructions for the type of equipment used. Calibrate equipment to
drop a teaspoon quantity (4 gams) of bail at 4-5 bot intervals in the artificial burrow made by the machine. Apply
at a rate of 2-3 lbs. of bail per acre. Pick up and burry all spitled bair.

ALFALFA

ALFALFA

ALFALFA

LOSE RESTRICTIONS: For control of meadow vales, Inorplaids voles, California voles. Degan voles, mountain voles and Terement's vales (Morota spe) and Fichardstein ground squares (Spormoghikas inclusiones) in a shalfa. All applications nut court and right and a califort of the law prote prote for the and spenith attailing a length of 2 notion. Adults loaged from treated assess must not be inversed until it reaches mature; This use is estimated to Merinac, California, factor, care del protection of the state of the control of the california of the ca

PREBATING: When bait is to be used to control Richardson's ground squirrels, prebaiting with untreated steamed crimped oats at a rate of 6 lbs, per acre 2-3 days before ZP® RODENT OAT BAIT AG™ is applied is likely to enhance acceptance of the toxic balt.

RPOADCAST RAITING: This product may be broadcast by air or province driven dispansion devices. Apply at a RIBADCAST BAITING: This product may be broadcast by air or ground-driven dispersing electors, Aprly all as the object to Riba, Class chie ingredient pare are. Make only not application per year if Richardson's ground qualities have being targeted. If voles are being targeted, a second balt application may be made after 10 days. A similarium of 2019. But, 6th a davie ingrediently are are may be applied per year. Do not apply, by air when wind aboutly exceeds 10 mph, Do not apply in piles or permit piles to be formed by equipment.

TIMOTHY AND TIMOTHY/ALFALFA MIXTURES PRODUCED FOR HAY USE RESTRICTIONS: For control of meadow votes, long-tailed votes, California votes, Oregon votes, mo

voles, and Townsend's voles (Microtus spp.) in timothy and timothy/alfalfa mixtures. Do not apply by air. Do not apply to actively growing timothy or timothy/alfalfa mixtures. This use is restricted to California, Idaho, Oregon and noton. Do not apply to timothy or timothy/alfalfa mixtures within 60 days of harvest,

Intersection or interruption of the property o

TIMOTHY PRODUCED FOR SEED

USE ESTIFICIONS For control of measure was to present the control of the control of measure makes, togetable values, california values. Output not mental make and himsensor beautiful of measure was togetable values. The control of the control of

devices at rates of up to 10 lbs. (0.2 lb, active ingredient) per acre during crop dominancy. Wear rubber gloves during hand-broadcast operations. A maximum of 20 lbs. (0.4 lb, active ingredient) per acre may be applied

USE RESTRICTIONS: For control of meadow voles, forcy allied voles, California voles, Conquired some voles, Monatenia voles, Consend's voles (Microtas sop.) and Pichardson's ground squirrieds (Spermophilis richardsonii) in growing-season and preplant applications in barley, Do not apply this product to barley within 50 days of harvest. This use is restricted to Montana, Idahin, Oregon and the state of Washington.

PREBAITING: When bait is used to control Richardson's ground squirrels, prebaiting with untreated steamed crimped dats at a rate of 6 bs, per acre 2-3 days before ZP® RODENT OAT BAIT AG™ is applied is likely to

enhance acceptance of the facts but in BRONCACST BATHER. This product may be broadcast by air or ground-rivinen dispensing devices. Apply ball at a rule up to B bb., 00, 12 bb. active ingredient) per acre. Make only one application per year if Richardsoris good sprained sending targeteds. It voles are been tageted, a score bad that application may be produced to the product of the product

DOTATORS

USE RESTRICTIONS: For control of meadow votes, long-tailed votes, California votes, Oregon votes, mountain voles, and Townsend's voles (Mcrotus spp.) in growing-season and pre-plant applications in potatoes. Do not apply this product to potatoes within 30 days of harvest. This use is restricted to Idaho, Oregon and the state of

BROADCAST BAITING: This product may be broadcast by air or ground-driven dispensing devices. Apply bait once at a rate up to 10 fbs. (0.2 fb, active ingredient) per acre, or make two separate applications at rates of up to 5 fbs. (0.1 fb, active ingredient) per acre. The maximum application rate per year is 10 fbs. (0.2 fb, active

SUGAR BEETS

USE RESTRICTIONS: For control of meadow votes, long-tailed votes, California votes, Oregon votes, mountain votes and Townsend's votes (Microtus sop.) in growing-season and preplant applications in supar beets, Do not apply this product to sugar beets within 30 days of harvest. This use is restricted to Idaho. Oregon and the state

BROADCAST BATTRG: This product may be broadcast by air or ground-driven dispensing devices, Apply bair at rates up to 10 bs, (0,2 bs, active ingredient) per acre at planting. A second application may be made at the same rate. The maximum application rate per year is 20 bs, (0,4 lb, active ingredient).

USE RESTRICTIONS: For control of meadow votes, broy-lailed votes, California votes, Congon votes, mountain votes, Tomsend's votes (Microtus spp.) and Richardson's ground squimels (Spermophilas inchardsonii) in growing-eason and replant applications in heate. Do not apply this product to wheat within 50 days of harvest, This use is restricted to Montana, Idaho, Oregon and the state of Washington.

PREBAITING: When bait is used to control Richardson's ground squirrels, prebaiting with untreated steamed crimped cats at a rate of 6 lbs. per acre 2-3 days before ZP® RODENT OAT BAIT AG™ is applied is likely to

RROADCAST RAITING: This product may be hyparcast by air or ground-driven dispension devices. Apply half an order of the first (1,12 ks, active) ingredient per are. Make only or application per year if Richardson's ground squirrels are being targeted. If voles are being targeted, a second application per year if Richardson's days, The maximum application rate per year is 12 bs. (0,24 b, active ingredient) per acre.

BEANS (DRY)
USE RESTRICTIONS: For control of prairie voles (Microtus ochrogaster), meadow voles (M. pennsylvanicus), house mice (Mus muscalus), deer mice (Percomyscus maniculatus), white-footed mice (P. feucopus), and Ord's kangaroo rats (Dipodomys ordil), in dry beans during the growing season. Do not apply by air. Do not apply this product to dry beans within 30 days of harvest. This use is restricted to Idaho, Oregon and the state of

BROADCAST BAITING: This product may be broadcast by hand or ground-driven dispensing devices at a rate

BUSHBERRIES AND CANEBERRIES

USE RESTRICTIONS: For control of meadow voles, prairie voles, and pine voles (Microtus spp.) in bushberries highbush and lowbush blueberries, currants, elderberries, gooseberries and huckleberries) and in caneberries lbtackberries, red raspberries, btack raspberries, loganberries, and cuttivars or hybrids of these caneberries). Only poly this product in the dormant season: after final harvest and not later than the beginning of leaf emerthe spring. Do not apply when ground is snow covered. Do not apply by air. Minimum preha

BROADCAST RAITING: This product may be broadcast by cycline seeder, or by hand. Wear rubber ployes during hand-broadcast operations. When applying by hand, throw tablespoon amounts (12 grams) into hecover along bushes, rock outcrops, and lence lines, Make up to 2 applications at a minimum interval of 21 de at the rate of 6 to 10 lbs, per acree per application.

CUCURRITS REDUCED THE AGE AND NO. THE CORN.

USE RESTRICTIONS: For control of prairie voles (Microtis ochrogassier), meadow voles (M. pennsylvanicus), house mice (Mis musculus), deer mice (Percomyacus manicualius) white-booled mice (P. Ruccopus), thirisen-fined ground squirrets (Spermophilus tridecemineatus), Ord's kangaroo rats (Dipodomys ordi), banner-tailed kangaroo s (D. spectabilis), Merriam's kangaroo rats (D. merriam), and cotton rats (Sigmodon hispidus) in reduced tillage and no-fill corn and in cucurbits, including and limited to chavate. Chinese waxoourd, citron melon, cucumbe gherkin, edible gourd. Momordica soo., muskmelon, pumpkin, squash (summer and winter), and watermelon. Only -furrow treatments are permitted for these crops. Do not broadcast bail. Do not apply this product within the

coupled habitate of the endingered species or subspecies of Nangaroo rats.

IN-FURROW APPLICATION ONLY: Equipment designed for in-furrow treatments must be used to apply this product in cucurbits, reduced tillage and no-fill corn, Make one application pre-planting, or at-planting, at a rate of 4 to 6 lbs, of balf per acre. Collect and remove excessive quantities of balf deposited on surface soil or crop ues during spills or equipment malfunctions.

CALIBRATION/ROW

Row spaces Inches	4 lbs./ A Rate oz. Per 1000 ft	5 lbs. / A rate oz. Per 1000 ft	6 lbs. / A rate oz. Per 1000 ft
20	2.5	3.1	3.7
30	3.7	4,6	5.5
36	4.4	5,5	6,6
38	4.7	58	69

TAWNS ORNAMENTALS COLE COLLEGE DADKS AND MIDSEDIES USE RESTRICTIONS: For control of meadow votes, prairie votes, pine votes, California votes, long-tailed votes, Oregon voles, mountain voles and Townsend's voles (Microtus spp.), Columbian ground squirrels (Spermophilus bianus). Richardson's ground squirrels (S. richardsoni), thirteen-lined ground squirrels (S. tridecerm(neatus), and California ground squirrels (S. beechevi) in lawns, ornamentals, golf courses, parks, and nurseries. Bait must not be applied on roads, over water or where plants are grown for food or feed,

PREBAITING: Pretoal by placing one teaspoon (4 grams) of untreated steamed crimped oats around each active burrow opening or in each active runway 2 or 3 days prior to using ZP® RODENT OAT BAIT AG™ to enhance

acceptance by ground squirrels or votes. HAND BATTING: Place 4 grams (one teaspoon) of bait around each active burrow opening or in each active nurway. Allow bait to fall through to the ground surface but do not apply bait to bare ground. Do not place bait in

USE RESTRICTIONS: This product may be used to control meadow votes (Microtus pennsylvanicus), prairie votes (M. cotrogaster), California votes (M. californicus), mountain votes (M. montanus), and prine votes (M. pinetis) around (within 15 feet of) agricultural, commercial and industrial buildings. Do not apply balt inside of structures. Bait must be placed in tamper-resistant bait stations. These stations must be resistant to destruction by doos and by children under 6 years of age and must be used in a manner that prevents such children from reaching into balt compartments and obtaining hait. If hait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger balt stations are needed in areas open to hoofed livestock, raccoons bears or other potentially destructive animals, or in areas prone to vandalism.

SELECTION OF TREATMENT AREAS: Determine where voles will most likely find and consume the bait, Look

for runways near buildings as well as signs of vole activity oriented toward the structures. PREBAITING: Prebaiting with untreated steamed crimped gats may improve accentance of this product and receptable of the scheduled application of 2P® RODENT OAT BAIT AG[™]. Check bait stations for evidence of bait consumption. Delay use of treated bait until more than half of the prebait offered has been consumed, move remaining prebait from bait stations before loading them with toxic bait.

BATTING: Apply a teaspoon (4 grams) of hait in each hait station. Space hait stations at intervals of 8-12 feet in tain balt in stations for 3 days. Extend the balting period only if there is co

AIRPORTS

USE RESTRICTIONS: This product may be used at airports to control meadow votes (Microtus pennsylvanicus), prairie voles (M. ochrogaster), California voles (M. californicus), mountain voles (M. montanus), pine voles (M. pinetis), and white-footed mice (Peromyscus leucopus). Do not apply bait to bare ground. Do not apply bait by air presses, are wine-nouses mick (retempscus excopuls), to not apply part to pare ground. Up not apply bast by air PREBATING: Prebailing with untreated steamed crimped cas may improve acceptance of this product. Apply prebait at the same rate and using the same procedures that are to be used for applying the toxic bait. Apply prebait 2 or 3 days before the scheduled application of ZP® RODENT OAT BAIT AG[™].

HAND BAITING: Place a teaspoon (4 grams) of bait in surface runways or at the mouths of holes leading to

mounted on a tractor or 4-wheel drive all-terrain vehicle (ATV). Wear rubber gloves during hand-broadcast operations. Apply treated bait at rates from 6 to 10 lbs, per acre. Do not broadcast bait on any area at an airport

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal,

STORAGE: Store only in original container in a cool, dry, well vertilated place inaccessible to children, pets and
wildlife. Protect from direct contact with water or excessive moisture. Once opened, do not store bag for prolonged

Spills: If container is punctured, repair with tape and use clear oversity bag and securely close. PESTICPE DEPOSAL: Pesticide wastles are toxic, Improper disposal of excess pesticide, spray minure, or insast is a violation of Federal Law If these wastes cannot be disposed of by use according to label instructions, contact, your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance,

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if

Dispose of empty container by placing in trash, at an approved waste disposal facility or by WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buver assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions

AG5081

040119/10-09