

SYNERGIZER®

Plant Food for Foliar and Soil Application

0-0-10

Product Information

SYNERGIZER 0-0-10 is a liquid plant food designed for foliar and soil application. **SYNERGIZER 0-0-10** is compatible with most fertilizer solutions and pesticides. Dilute **SYNERGIZER 0-0-10** with water before mixing with other fertilizers or pesticides.

GUARANTEED ANALYSIS

Soluble Potash (K ₂ O).....	10%
Magnesium (Mg).....	0.3%
Iron (Fe).....	1%
1.0% Chelated Iron	
Manganese (Mn).....	0.2%
Zinc (Zn).....	0.5%

Derived from potassium hydroxide, magnesium oxide, manganese oxide, ferric ammonium citrate, and zinc oxide.

F1416

KEEP OUT OF REACH OF CHILDREN

**SHAKE WELL BEFORE USE
DO NOT APPLY AT TIME OF BLOOM**



WARNING

Harmful if swallowed.

Precautionary statement: Avoid contact with eyes. **First Aid:** if swallowed: seek medical attention. Rinse mouth. If in eyes: remove contact lenses if present. Wash thoroughly with water for several minutes. **Storage & Disposal:** store in a safe manner and away from incompatible materials. Dispose product/container in accordance with federal, state, and local regulations.

NET CONTENTS: 2.5 gal. (9.45 l)
Weight per Gallon: 10.5 lb. (4.7 kg) @ 68°F
Density: 1.2g/ml @ 68°F
Lot #:

**Made in
USA**

 **JHBiotech, Inc.**
Innovation for a Greener Earth

MANUFACTURED BY: JH BIOTECH, INC.
P.O. Box 3538, Ventura, CA 93006 U.S.A.
805-650-8933 jhbiotech.com

Directions and General Recommendations

Generally, dilute **SYNERGIZER 0-0-10** with at least 50 times of water. Apply to plants according to the following recommendations:

ALFALFA: 2-4 qt./ac. one week after each cutting.

ALMONDS, APRICOTS, CHERRIES, HOPS, PLUMS, PEACHES: 3-4 qt./ac. after bud swell to popcorn stage, prior to bloom. Older trees may require another application 4-6 weeks later.

BEANS, PEAS: 2-3 qt./ac. at bud stage, prior to bloom. Repeat the application 2 more times at 2 week intervals.

CITRUS: 4-6 qt./ac. at bud stage, prior to bloom. When new growth appears, repeat the application 2 more times at 2-4 week intervals.

COTTON: 1-2 qt./ac. after true leaves appear. Second application should be made at early square at 2-3 qt./ac. Repeat the application 2 more times at 2 week intervals.

POTATO: 1-2 qt./ac. at 6-8 leaf stage. 1-3 qt./ac. at tuber initiation. 2-3 qt./ac. when tubers are 1-2 in. diameter. Repeat the application 2 more times at 1-2 week intervals.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

STRAWBERRY: 2-4 qt. starting after plants are established. Repeat the application at 2-3 week intervals throughout the season.

TOMATO AND PEPPER: 1-2 qt./ac. after stand is established. 1-3 qt./ac. at bud stage, prior to bloom. Repeat the application at 1-3 qt./ac. at 2 week intervals.

TURF: Apply 2-3 fl. oz. per 1,000 sq. ft. in 4-10 gal. of water. Apply as needed.

SYNERGIZER 0-0-10 can be used on many crops including but not limited to alfalfa, almonds, apple, apricots, avocado, banana, barley, beans, carrot, cherries, citrus, corn, cotton, grapes, hops, mango, ornamental crops, palm, papaya, peaches, pear, pecan, pepper, pistachio, plantain, plums, potato, prune, rice, row crops, sorghum, soybean, strawberry, sugarbeet, sunflower, tomato, turf, vegetables, walnut, and wheat.

059004-D-B-1

Copyright © 2018 All rights reserved